arXiv:1511.01003v1 [astro-ph.HE] 3 Nov 2015

Luminosity of ultra high energy cosmic rays and bounds on magnetic luminosity of radio-loud AGNs

C. H. Coimbra-Araújo* and R. C. Anjos†
Departamento de Engenharias e Ciências Exatas, Universidade Federal do Paraná (UFPR), Pioneiro, 2153, 85950-000 Palotina, PR, Brazil.

We investigate the production of magnetic flux from rotating black holes in active galactic nuclei (AGNs) and compare it with the upper limit of ultra high energy cosmic ray (UHECR) luminosities, calculated from observed integral flux of GeV-TeV gamma rays for nine UHECR AGN sources. We find that, for the expected range of black hole rotations ($0.44 < a < 0.80$), the corresponding bounds of theoretical magnetic luminosities from AGNs coincides with the calculated UHECR luminosity. We argue that such result possibly can contribute to constrain AGN magnetic and dynamic properties as phenomenological tools to explain the requisite conditions to proper accelerate the highest energy cosmic rays.

PACS numbers: 13.85.Tp, 98.70.Sa, 95.30.Qd, 98.62.Js

The origin of the highest energy cosmic rays, or ultra high energy cosmic rays (UHECRs), with energies $E > 10^{18}$ eV, in spite of huge observational/experimental endeavors, represents one of the greatest puzzles of modern astrophysics [1, 2]. The possibility of accelerating particles up to such extreme energies is addressed by the well-known Hillas source plot [3]. There, a table of possible UHECR sources is presented by bringing the simple yet efficient remark that particles, during acceleration, are confined in the source on a Larmor timescale (for more details see, e.g., [1, 2, 4–9]).

The prominent extragalactic candidates for accelerating particles (mainly protons) to the highest energies are active galactic nuclei (AGNs) [10], the most powerful radiogalaxies [11, 12], and also gamma ray bursts, fast spinning newborn pulsars, interacting galaxies, large-scale structure formation shocks and some other objects [1].

On the other hand, the reconstruction of cosmic ray luminosities, from Earth laboratory observations, possibly can shed some light on radiative bounds of UHECR potential sources. For example, in [13, 14] it is shown that using the methods of UHECR propagation from the source to Earth and the measured upper limit on the integral flux of GeV-TeV gamma-rays it is possible to infer the upper limits of the proton and total UHECR (iron) luminosity. This comes from the fact that gamma-rays can be produced as a result of the cosmic ray propagation and contribute to the total flux measured from the source.

In the present work, some of the potential AGN sources will be investigated as UHECR sources (see Table I). It will be calculated upper limits of UHECRs luminosities to be compared to the theoretical magnetic/jet luminosity of those AGNs. To reconstruct, from experiments, the possible UHECR luminosities here it will be used the method described in [13] as a prolific way to calculate upper limits on UHECR luminosities.

In first place, space and ground instruments, as FERMI-LAT [15], VERITAS [16], H.E.S.S. [17] and MAGIC [18] provide upper limits on the GeV-TeV gamma-ray integral flux and the method [13] connects those measured upper limits with the source UHECR cosmic ray luminosity (L_{CR}^{UL}) by

$$L_{CR}^{UL} = \frac{4\pi D_s^2 (1+z_s)\langle E\rangle_0}{K_\gamma \int_{E_{th}}^{\infty} dE_\gamma \; P_\gamma(E_\gamma)} \; I_\gamma^{UL}(> E_\gamma^{th}), \tag{1}$$

where $I_\gamma^{UL}(> E_\gamma^{th})$ is the upper limit on the integral gamma-ray flux for a given confidence level and energy threshold, K_γ is the number of gamma rays generated from the cosmic ray particles, $P_\gamma(E_\gamma)$ is the energy distribution of the gamma-rays arriving on Earth, E_γ is the energy of gamma-rays, $\langle E\rangle_0$ is the mean energy, D_s is the comoving distance and z_s is the redshift of the source. This method allows one to calculate upper limits on the proton and total luminosities for energies above 10^{18} eV. Also, it illustrates techniques to study the origin of UHECRs from multi-messenger GeV-TeV gamma-rays and it has been used to calculate at least upper limits for thirty sources (AGNs), with redshift smaller than 0.048 and UHECR spectra measured by the Pierre Auger [19] and Telescope Array [20] (TA) observatories. In fact, as described by [13, 14], the UHECR upper limit luminosity is obtained from the integral of the gamma-ray flux from the observed spectrum of UHECRs. Propagation models and the measured upper limit of gamma-ray flux of a source amongst propagation models are the fundamental aspects to be considered to perform

*Electronic address: carlos.coimbra@ufpr.br
†Electronic address: ritacassia@ufpr.br

such calculation. Indeed, this same method will allow, in the future, from CTA Observatory [21], a range of new UHECR luminosity upper limits.

In second place, it is well-known that radiative mechanisms within AGNs are chiefly powered by their central supermassive black holes (SMBHs) and the companion accretion disk. The rotation of the system creates twisted magnetic fields that drive jets of relativistic particles. Here it will be considered a standard accretion mechanism (Bondi accretion model $\dot{M} = \pi\lambda c_S \rho_B r_B$ [22], where $\dot{M}$ is the accretion rate, λ takes the value of 0.25 for an adiabatic index 5/3, c_S is the sound speed in the medium, r_B is the Bondi accretion radius and ρ_B is the gas density at that radius) which produces, by friction and other radiative processes, enormous bolometric luminosities. The injection of only a moderate fraction of this bolometric luminosity would suffice to reproduce the observed cosmic ray flux above 10^{19} eV. Nevertheless, it appears that cosmic ray flux from AGNs probably comes from jet luminosities. Therefore, the magnetic field of the structure is the main responsible to produce such perpendicular jet outflows. In this case, they do not all offer as appetizing physical conditions with respect to particle acceleration as, e.g., gamma ray bursts, since considerable outflow luminosities, i.e., magnetic luminosities L_B are actually required to accelerate protons to the highest energies observed. This limits the potential number of AGNs as cosmic ray sources in the nearby universe (unless the highest energy cosmic rays are heavy nuclei) [2]. In the present contribution, we describe the mechanism via powering jets from Blandford-Znajek mechanism [23]. This approach is based on flux accumulation that leads to the formation of a magnetically chocked accretion flow and strong flux of high energy particle jets and it assumes that any geometrically thick or hot inner region of an accretion flow can drive magnetic field fluctuations to produce jets. This technique has been proposed to explain the most luminous and radio-loud AGNs, and consequently could also explain mechanisms behind the production of UHECRs.

In what follows it is derived a description for the production of UHECR luminosities based on the possible relation between magnetic flux accumulation and jet production efficiency. Similar schemes have been proposed to describe UHECR luminosity contribution and sources of cosmic rays from black hole accretion mechanisms [24, 25]. The significant feature of this model is that the dominant factor in the magnetic luminosity is the powerful jets to therefore determinate the radio loudness of the AGN. In first place, considering a system with a rotating central BH, the necessary condition to use Blandford-Znajek mechanism is that $\Phi_d > \Phi_{BH,\max}(\dot{M})$, i. e., the net poloidal magnetic flux Φ_d trapped in the disk is larger than the maximum that can be confined on the BH caused by pressure of the accreting plasma. Satisfied this condition, the rate of energy extraction from the rotating BH via the Blandford-Znajek mechanism yields the magnetic luminosity

$$\begin{aligned} L_B &\simeq 4\times 10^{-3}\Phi^2_{BH,\max}(\dot{M})\frac{\Omega^2_{BH}}{c} f_a(\Omega_{BH}) \\ &= 10(\phi/50)x_a^2 f_a(x_a)\dot{M}c^2, \end{aligned} \tag{2}$$

where Ω_{BH} is the angular velocity of the black hole and

$$x_a = r_g\frac{\Omega_{BH}}{c} = [2(1+\sqrt{1-a^2})]^{-1}a, \tag{3}$$

with

$$f_a(x_a) \simeq 1 + 1.4x_a^2 - 9.2x_a^4, \tag{4}$$

where a is the dimensionless angular momentum parameter ($a = J/Mc$, with J the BH angular momentum), ϕ is a dimensionless factor which, according to numerical simulations (see, e.g., [26]), is typically of order 50, and r_g is the gravitational radius $r_g = GM/c^2$.

In Fig. 1 it is displayed the effect of black hole spinning on magnetic luminosity with the variation of mass. The greater the AGN BH spinning, the greater the AGN magnetic loudness. A similar result is observed with the increasing of the black hole mass. In this approach, M and a are therefore the main parameters to affect energetic jets and the consequent energetic cosmic ray production at AGN expected sources. For comparison one can calculate the upper limit of bolometric luminosity for each source, i.e., the Eddington luminosity $L_{\rm Edd} = \frac{4\pi GMm_pc}{\sigma_T} = 1.26\times 10^{45}\frac{M}{10^7 M_\odot}$ (where M is the mass of the AGN, m_p is the hydrogen atom mass, c is the light speed and σ_T is the Stephen-Boltzmann constant) to see that, as awaited, for each source $L_B < L_{\rm Edd}$.

As AGN jets can be the main extragalactic sources of UHECRs [10, 27], one can write the UHECR luminosity L_{CR}^{Theory} as a fraction η of the magnetic luminosity:

$$L_{CR}^{Theory} = \eta L_B. \tag{5}$$



FIG. 1: L_B as a function of the mass and parameter a. For all plots it was considered $\dot{M}$ as the Bondi accretion rate [22]. For comparison, the Eddington luminosities $L_{\rm Edd}$ for each considered AGN mass are: $L_{\rm Edd}(M = 10^{9.5} M_\odot) = 1.26 \times 10^{47.5}$ erg s^{-1}, $L_{\rm Edd}(M = 10^{10.0} M_\odot) = 1.26 \times 10^{48}$ erg s^{-1}, $L_{\rm Edd}(M = 10^{10.5} M_\odot) = 1.26 \times 10^{48.5}$ erg s^{-1}, $L_{\rm Edd}(M = 10^{11.0} M_\odot) = 1.26 \times 10^{49}$ erg s^{-1}.

For example, bounds on the fraction η_{pr} of L_B to be converted in relativistic protons, as a function of BH spin a, come from eqs. (??) and (??) as

$$\eta(a)_{pr} = \frac{L_{pr}^{UL}}{10(\phi/50)x_a^2 f_a(x_a)\dot{M}c^2}. \tag{6}$$

For the spin range $0.44 < a < 0.8$ where most black holes are expected to lie, the fraction η_{pr} varies from as $\sim 5\%$ to 40% for $a = 0.45$ and from 2% to 10% for $a = 0.8$, see Fig. 2. An important remark is that Fig. 2 reflects the outter bounds of all nine curves $\eta(a)_{pr}$ produced from the nine sources of Table I. The sources 2MASX J1145 and NGC 5995 are taken as the critical limits for AGNs in the universe up to redshifts $z_s < 0.048$ and they plot the threshold curves for eq. (??). See, for instance, that the upper limit L_{pr}^{UL} for these two sources is greater than the theoretical calculated range (for a spin $a \sim 0.7$). In this case, it is expected that their masses produce low levels of magnetic luminosity. To explain such unexpected great upper limits, one has to admit, e.g., that possibly those AGNs have a critical spinning black hole, with $a \to 1$, since the greater the spin, the greater the luminosity L_B.



FIG. 2: η_{pr} as a function of the parameter a. For all plots it was considered $\dot{M}$ as the Bondi accretion rate [22].

Table I shows the range $L_{CR,\rm min}^{Theory} - L_{CR,\rm max}^{Theory} = \eta_{pr,\rm min} L_B - \eta_{pr,\rm max} L_B$ for nine AGN sources, assuming a fixed spin of $a = 0.45$ and $a = 0.7$. The column L_{pr}^{UL} is the calculated upper limit of the proton luminosity for each source from eq. (??), and it has dependence only with observational [GeV-TeV gamma rays + cosmic rays] constraints

[13, 14], i.e., it has no dependence with BH mass or BH spin of the AGN. Comparing both results, it is possible to see that NGC 1142 fails to produce a proton upper luminosity that remains in the L_{CR}^{Theory} calculated range. This is explained by the fact that the observed NGC 1142 mass must drive a great magnetic luminosity L_B, since the greater the mass, the greater the luminosity L_B. To explain the low cosmic ray luminosity from NGC 1142 one can assume that the AGN black hole possibly has a small spin, since the smaller the spin, the smaller the luminosity L_B and consequently the smaller the cosmic ray luminosity.

The present work compared ultra high energetic cosmic ray luminosities with a theoretical luminosity derived from intrisic AGN properties, as its mass and central black hole spin. An important remark is that luminosity calculation from method (**??**) does not require any AGN property, since they come unically from observed integral flux of GeV-TeV gamma-rays of UHECR AGN sources. In this aspect, both for proton and iron luminosities it is possible to find phenomenological bounds on the conversion fraction of magnetic luminosities into energetic particles. Such results logically lead to the question whether AGNs are or not the sources of the high energy cosmic rays. In general, the Fermi acceleration processes are evoqued to explain if AGNs could indeed reach $\sim 10^{19.5}$ eV. In parallel, for protons, only the most powerful flat spectrum radio quasars, which are considered to be the jet-on analogs of FR II radio galaxies with relativistic jets, show a magnetic luminosity in excess of 10^{45} ergs s^{-1} that can power sufficient energetic jets. Those are the objects studied here. In complement, BL Lac objects or TeV blazars, thought to be the analogs of FR I radio galaxies (such as Cen A), typically exhibit magnetic luminosities L_B of the order of 10^{44} ergs s^{-1} or less. In such cases, only Fermi processes and shocks in the jets and the hot spots of the most powerful FRII radio-galaxies may nevertheless offer the requisite conditions to proper accelerations to reach, e. g., $\sim 10^{20}$ eV cosmic rays [11, 12, 28]. In this manner, the presented results could possibly contribute to constrain magnetic and dynamic properties of AGNs to better understand the requisite conditions to proper accelerate the highest energy cosmic rays.

Acknowledgments

The authors are very grateful to the researchers of DEE-UFPR. This work was funded by CNPq under grant 458896/2013-6.

[1] K. Kotera and A. V. Olinto, *The Astrophysics of Ultrahigh-Energy Cosmic Rays*, Ann. Rev. of Astron. and Astrophys. **49**, 119 (2011).
[2] M. Lemoine, *Acceleration and propagation of ultrahigh energy cosmic rays*, JPhCS **409**, 012007 (2012).
[3] A. M. Hillas, *The origin of ultra-high-energy cosmic rays*, Ann. Rev. Astron. Astrophys. **22**, 425 (1984).
[4] M. Nagano and A. A. Watson, *Observations and implications of the ultrahigh-energy cosmic rays*, Rev. Mod. Phys. **72**, 689 (2000).
[5] M. Nagano, *Search for the end of the energy spectrum of primary cosmic rays*, N. J. of Phys. **11**, 065012 (2009).
[6] J. Bluemer, R. Engel, J. R. Hörandel, *Cosmic rays from the knee to the highest energies*, Prog. in Part. and Nuc. Phys. **63**, 293 (2009).
[7] J. Beatty and S. Westerhoff, *The Highest-Energy Cosmic Rays*, Annu. Rev. Nucl.Sci and Particle Physics **59**, 319 (2009).
[8] Letessier-Selvon, A and T Stanev, *Ultrahigh energy cosmic rays*, Rev. Mod. Phys. **83**, 907 (2011).
[9] A. A. Watson, *High-energy cosmic rays and the GreisenâĂŞZatsepinâĂŞKuz'min effect*, Rep. Prog. Phys. **77**, 036901 (2014).
[10] G. Henri, G. Pelletier, P.-O. Petrucci, N. Renaud, *Active galactic nuclei as high energy engines*, Astropart. Phys. **11**, 347 (1999).
[11] F. Takahara, *On the origin of highest energy cosmic rays*, Prog. Theor Phys. **83**, 1071 (1990).
[12] J. P. Rachen, P. L. Biermann, *Extragalactic ultra-high energy cosmic-rays-part one-contribution from hot spots in Fr-II radio galaxies*, Astron. Astrophys. **272** 161 (1993).
[13] A.D. Supanitsky and V. de Souza, *An upper limit on the cosmic-ray luminosity of individual sources from gamma-ray observations*, *JCAP*, **12**, 023 (2013).
[14] R. C. Anjos, A.D. Supanitsky and V. de Souza, *Upper limits on the total cosmic-ray luminosity of individual sources* , *JCAP*, **07**, 049 (2014).
[15] The Fermi-LAT Collaboration, *The first catalog of active galactic nuclei detected by the Fermi Large Area Telescope*, *The Astrophysical Journal*, **715**, 429 (2010).
[16] N. Galante, for the VERITAS Collaboration, *VERITAS Observations of Radio Galaxies*, *Fermi Symposium* (2009) 2 arXiv:0912.3850.
[17] F. Aharonian et al., *Upper limits from HESS active galactic nuclei observations in 2005-2007*, *Astronomy & Astrophysics*, **478**, 387 (2008).
[18] The MAGIC Collaboration, *Detection of very high energy gamma-ray emission from the Perseus cluster head-tail galaxy IC 310 by the MAGIC telescopes* , (2010), arXiv:1009.2155.

[19] A. Schulz for The Pierre Auger Collaboration, *The measurement of the cosmic ray spectrum above* 3×10^{17} *eV with the Pierre Auger Observatory*, *Proceedings of the 33rd International Cosmic Ray Conference*, Rio de Janeiro, Brazil, 2013, arXiv:1307.5059.

[20] Douglas Bergman for the Telescope Array Collaboration, *TA Spectrum Summary*, *Proceedings of the 33rd International Cosmic Ray Conference, Rio de Janeiro, Brazil, 2013.*

[21] CTA Consortium,*Design Concepts for the Cherenkov Telescope Array*, *Experimental Astronomy*, **32** (2011) 193-316.

[22] H. Bondi, *On Spherically Symmetrical Accretion*, *MNRAS* **112**, 195 (1952).

[23] M. Sikora and M. C. Begelman, *Magnetic flux paradigm for radio-loudness of AGN*, *APJ* **64**, L24 (2013).

[24] I. Dutan, Laurentiu I. and I. Caramete, *Ultra-high-energy cosmic rays from low-luminosity active galactic nuclei*, *Astro. Phys.*, **62** (2015) 206.

[25] J.K. Becker and P.L. Biermann, *Neutrinos from active black holes, sources of ultra high energy cosmic rays*, *Astro. Phys.*, **31** (2009) 138.

[26] J. C. McKinney, A. Tchekhovskoy, R. D. Blandford, *General relativistic magnetohydrodynamic simulations of magnetically choked accretion flows around black hole*, MNRAS **423**, 3083 (2012).

[27] P. L. Biermann and P. A. Strittmatter, *Synchrotron emission from shock waves in active galactic nuclei*, *Astrophys. J.* **322**, 643, 1987.

[28] A. Celotti and G. Ghisellini, *The power of blazar jets*, Month. Not. Roy. Astron. Soc. **385**, 283 (2008).

[29] E. Waxman and J. Bahcall, Phys. Rev. Lett. **78**, 2292 (1997).

[30] E. Waxman, Lect. Notes in Phys. 576, **122** (2001).

[31] R. Abbasi et al. (The Ice Cube Collaboration), Nature **484**, 351 (2012).

[32] Lisa M.W., *Extragalactic X-ray Surveys of ULXs and AGNs*, dissertation, University of Maryland, 2008.

[33] Michael Koss et al, *Host Galaxy Properties of the Swift Bat Ultra Hard X-Ray Selected Active Galactic Nucleus*, ApJ **739** 57 (2011).

[34] K. Oh, H. Choi, H.-G. Kim, J.-S. Moon, S. K. Yi, Astron. J. **146**, 151O (2013).

TABLE I: Comparison between cosmic ray luminosity L_{pr}^{UL} (protons) from the method derived from eq. (**??**) and the theoretically calculated range $L_{CRmin}^{Theory} - L_{CRmax}^{Theory}$ of cosmic ray luminosities to $a = 0.45$ and $a = 0.7$, from many sources. For comparison, it is also calculated the upper limit of the bolometric luminosity (the Eddington luminosity $L_{\rm Edd}$) for each case below, in erg s^{-1}: $L_{\rm Edd}[\rm NGC985] = 5.02 \times 10^{48}$, $L_{\rm Edd}[\rm NGC1142] = 6.31 \times 10^{48}$, $L_{\rm Edd}[\rm 2MASXJ07] = 4.68 \times 10^{48}$, $L_{\rm Edd}[\rm CGCG420] = 6.83 \times 10^{48}$, $L_{\rm Edd}[\rm MCG-01] = 1.82 \times 10^{48}$, $L_{\rm Edd}[\rm 2MASXJ11] = 1.26 \times 10^{48}$, $L_{\rm Edd}[\rm LEDA] = 4.92 \times 10^{48}$, $L_{\rm Edd}[\rm NGC5995] = 9.78 \times 10^{47}$, $L_{\rm Edd}[\rm Mrk520] = 3.17 \times 10^{48}$. The mass source comes from [32, 33]. Redshift and other properties are taken from [34].

Source name	z_s	$\log M_\odot$	$\mathbf{L_{pr}^{UL}}$ (Proton) [erg s^{-1} $\times 10^{45}$]	$\mathbf{L_B}$[erg s^{-1} $\times 10^{45}$] $a = 0.45$	$a = 0.7$	$a = 0.45$ $\mathbf{L_{CRmin}^{Theory} - L_{CRmax}^{Theory}}$ [erg s^{-1} $\times 10^{45}$]	$a = 0.7$ $\mathbf{L_{CRmin}^{Theory} - L_{CRmax}^{Theory}}$ [erg s^{-1} $\times 10^{45}$]
NGC 985	0.04353	10.6	1.03	5.14	15.54	0.21 - 2.57	0.77 - 2.79
NGC 1142	0.02916	10.7	0.49	6.47	19.57	0.26 - 3.23	0.97 - 3.52
2MASX J07595347+2323241	0.03064	10.57	1.01	4.80	14.51	0.19 - 2.40	0.72 - 2.61
CGCG 420-015	0.02995	10.63	0.95	5.51	16.66	0.22 - 2.76	0.83 - 2.99
MCG-01-24-012	0.02136	10.16	0.65	1.86	5.64	0.07 - 0.93	0.28 - 1.02
2MASX J11454045-1827149	0.03616	10.0	1.30	1.29	3.90	0.07 - 0.64	0.19 - 0.70
LEDA 170194	0.04024	10.59	1.48	5.02	15.19	0.21 - 2.51	0.75 - 2.73
NGC 5995	0.02834	9.89	0.90	1.00	3.03	0.04 - 0.51	0.15 - 0.54
Mrk 520	0.02772	10.4	0.98	3.24	9.81	0.13 - 1.62	0.49 - 1.77